

Modeling finite entropy states with free fermions

Oleksandr Gamayun

University of Amsterdam & BITP

arXiv:2012.02079 +
work-in-progress together with:


N. Iorgov and Yu. Zhuravlev & D. M. Chernowitz and J.S. Caux

11/02/2021
LIJC (Zoom)

Correlation functions in 1D systems

$$\langle \mathbf{q} | \mathcal{O}(x, t) \mathcal{O}(0, 0) | \mathbf{q} \rangle = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + ixP_{\mathbf{k}}}$$

- ▶ Numerics (ABACUS)
- ▶ Field theory ($k_F x \gg 1$, $k_F^2 t \gg 1$)
 - ▶ Linear Spectrum
 - ▶ $\mathcal{O} = P(\partial\varphi, \partial^2\varphi, e^{i\varphi})$
 - ▶ Universality from microscopic




Left: Comparison between ABACUS and inelastic neutron scattering for $KCuF_3$. [PRL 111 137205].
 Right: The threshold singularities in the Non-Linear Luttinger Liquid.

Correlation functions in 1D systems

$$\langle \mathbf{q} | \mathcal{O}(x, t) \mathcal{O}(0, 0) | \mathbf{q} \rangle = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + ixP_{\mathbf{k}}}$$

- ▶ Numerics (ABACUS)
- ▶ Field theory ($k_F x \gg 1$, $k_F^2 t \gg 1$)
 - ▶ Linear Spectrum
 - ▶ $\mathcal{O} = P(\partial\varphi, \partial^2\varphi, e^{i\varphi})$
 - ▶ Universality from microscopic



Left: Comparison between ABACUS and inelastic neutron scattering for $KCuF_3$. [PRL 111 137205].
 Right: The threshold singularities in the Non-Linear Luttinger Liquid.

Both approaches fail at finite temperature !!!

Outline

- ▶ Observables
- ▶ Sine-kernel
 - ▶ Microscopic bosonization $T = 0$
 - ▶ Phase dressing
 - ▶ Static correlations
- ▶ Lattice Sine-kernel (XY-model)
 - ▶ Effective form-factors
 - ▶ Summation
 - ▶ From Fredholm to Toeplitz
- ▶ Dynamics

Observables

$$\tau(x) = \det(1 + \hat{V})$$



$$\hat{V}f(q) = \int_{\gamma} dq' V(q, q') f(q')$$

- ▶ Effective numerical evaluation (F. Bornemann "On the Numerical Evaluation of Fredholm Determinants" [0804.2543])

$$\int_a^b f(q) dq = \lim_{N \rightarrow \infty} \sum_{k=1}^N \omega_k f(x_k), \quad \det(1 + \hat{K}) = \lim_{N \rightarrow \infty} \det(\delta_{ij} + \sqrt{\omega_i} K(x_i, x_k) \sqrt{\omega_k}) \Big|_{1 \leq i, k \leq N}$$

- ▶ Finite entropy state

$$V(p, q) \rightarrow n_F(p) V(p, q)$$

Generalized Sine Kernels

$$\tau(x) = \det \left(1 + \frac{e^{2\pi i\nu} - 1}{\pi} n_F(q) \frac{\sin x(q-p)/2}{q-p} \right)$$

- ▶ Random matrices
- ▶ Impenetrable bosons
- ▶ Mobile impurity [SciPost Phys. 8, 053 (2020), New J. Phys. 18 (2016), 045005]

$$\rho(y) = \det(1 + \hat{K} + \delta\hat{K}) - \det(1 + \hat{K})$$

- ▶ Return probability from the domain wall initial state $|\text{DW}\rangle = |\uparrow\uparrow \dots \uparrow\downarrow \dots \downarrow\rangle$ [J.M. Stephan, J. Stat (2017)]

$$\langle \text{DW} | e^{\tau H_{\text{XXX}}} | \text{DW} \rangle = \det_{\mathbb{R}_+} \left(1 - e^{-\rho^2/4} \frac{\sin \sqrt{\tau}(\rho - q)}{\pi(\rho - q)} e^{-q^2/4} \right)$$

- ▶ Persistence of spin configurations [I. Dornic (2018)] $n_F(q) = 1/\cosh(q)$
- ▶ Classical integrable systems $n_F(q) = r(q)$

Generalized Sine Kernels

$$\tau(x) = \det_{[-\pi, \pi]} \left(1 + \hat{V} + \delta \hat{V} \right) - \det_{[-\pi, \pi]} \left(1 + \hat{V} \right)$$

XY model [A.G. Izergin, V.S. Kapitonov, N.A. Kitanine, solv-int/9710028]

$$V(p, q) = -\frac{\omega_F(q)}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}}, \quad \delta V(p, q) = -\frac{\omega_F(q)}{\pi} e^{-i(p+q)x/2} e^{-\frac{i(p-q)}{2}}$$

$$\omega_F(q) = \frac{1}{2} \left(1 - e^{i\theta(q)} \tanh \frac{\beta E(q)}{2} \right)$$

$$\mathbf{H}_{XY} = -\frac{1}{2} \sum_{j=1}^L \left[\frac{1+\gamma}{2} \sigma_j^x \sigma_{j+1}^x + \frac{1-\gamma}{2} \sigma_j^y \sigma_{j+1}^y + h \sigma_j^z \right], \quad \tau(x) \equiv \frac{\text{Tr} \sigma_{x+1}^x \sigma_1^x e^{-\beta \mathbf{H}_{XY}}}{\text{Tr} e^{-\beta \mathbf{H}_{XY}}}$$

Spectrum of fermionic (Majorana) excitations

$$E(q) = \sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}$$

Bogolyubov rotation angle

$$e^{i\theta(q)} = \frac{h - \cos q - i\gamma \sin q}{\sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}}$$

Sine Kernel ($T = 0$)

$$\tau(x) = \tau(x, t = 0) = \det_{[-k_F, k_F]} \left(1 + \frac{e^{2\pi i\nu} - 1}{\pi} \frac{\sin x(q - p)/2}{q - p} \right)$$

Form-factor presentation

$$\tau(x, t) = \langle \mathcal{O}(x, t) \mathcal{O}(0, 0) \rangle = \sum_{k_1 < k_2 < \dots < k_N} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + iP_{\mathbf{k}}x}$$

$$|\mathbf{q}\rangle: \text{free fermions: } q_i = \frac{2\pi n_i}{L}; \quad |\mathbf{k}\rangle: \text{shifted free fermions: } k_i = \frac{2\pi(n_i - \nu)}{L}$$

$$P_{\mathbf{k}} = \sum_i k_i, \quad E_{\mathbf{k}} = \sum_i k_i^2/2$$

The form-factor (overlap):

$$|\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 = \left(\frac{2}{L} \sin \pi\nu \right)^{2N} \left(\det_{N \times N} \frac{1}{k_i - q_j} \right)^2 \rightarrow |\langle \mathbf{q} | \mathbf{k} \rangle|^2.$$


Field theory treatment = Microscopic bosonization ($T = 0$)

- ▶ Form-Factor summation

$$\tau(x, t) = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}}} = \det(1 + \hat{V})$$

- ▶ Orthogonality Catastrophe: $|\langle \mathbf{q} | \mathcal{O} | \mathbf{k}_{\text{vac}} \rangle|^2 = \mathcal{A}/N^{2\alpha}$
- ▶ Soft-mode summation

$$\begin{aligned} \tau(x) &\sim \sum_{\text{IR}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}}} = \sum_{k=0}^{\infty} \sum_{\substack{p_1, \dots, p_k \\ h_1, \dots, h_k}} \langle \Omega | e^{\sqrt{\alpha}\varphi(x,t)} | \{p, h\} \rangle \langle \{p, h\} | e^{-\sqrt{\alpha}\varphi(0,0)} | \Omega \rangle \\ &= \mathcal{A} e^{-i(P_{\Omega} - P_{\text{vac}})x} \langle e^{\sqrt{\alpha}\varphi(x)} e^{-\sqrt{\alpha}\varphi(0)} \rangle = \frac{\mathcal{A}}{x^{2\alpha}} e^{-i(P_{\Omega} - P_{\text{vac}})x} \end{aligned}$$



Slavnov (1989); Slavnov and Korepin (1991); A. Shashi, L. I. Glazman, J.-S. Caux, and A. Imambekov (2011); N. Kitanine, K.K. Kozłowski, J.-M. Maillet, N.A. Slavnov, and V. Terras (2009-2012); K.K. Kozłowski, J.-M. Maillet (2015);

Combinatorics of orthogonality catastrophe

Generic overlap

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \left(\frac{2}{L} \sin \pi \nu \right)^{2N} \left(\det_{N \times N} \frac{1}{k_i - q_j} \right)^2 = \left(\frac{2}{L} \sin \pi \nu \right)^{2N} \frac{\prod_{i>j} (k_i - k_j)^2 \prod_{i>j} (q_i - q_j)^2}{\prod_{i,j} (k_i - q_j)^2}.$$

Fermi sea integers

$$k_j = \frac{2\pi}{L} (n_j - \nu), \quad q_j = \frac{2\pi}{L} n_j, \quad n_j = -\frac{N-1}{2} + j - 1, \quad j = 1, 2, \dots, N$$

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \left(\frac{\sin \pi \nu}{\pi \nu} \right)^{2N} \prod_{i \neq j} \left(1 - \frac{\nu}{i-j} \right)^{-2} = \frac{G^2(1-\nu)G^2(1+\nu)G^4(N+1)}{G^2(N-\nu+1)G^2(N+\nu+1)}.$$


$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \frac{G^2(1-\nu)G^2(1+\nu)}{N^{2\nu^2}}$$

For $\nu = \nu(k)$, ($\nu_{\pm} = \nu(\pm k_F)$), $k_F = \pi L/N$

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \frac{G^2(1-\nu_-)G^2(1+\nu_+)(2\pi)^{\nu_- - \nu_+}}{N^{\nu_-^2 + \nu_+^2}} \exp \left(\int_{[-k_F, k_F]^2} \frac{\nu(\lambda)\nu(\mu)}{(\lambda - \mu + i0)^2} d\lambda d\mu \right)$$

Static + zero temperature

$$\tau(x) = \frac{G^2(1-\nu)G^2(1+\nu)}{(-2ix)^\nu(2ix)^\nu} e^{-2i\nu x} + (\nu \rightarrow \nu + \mathbb{Z})$$



Finite temperature

$$\tau(x) = \det \left(1 + \frac{n_F(q)}{\pi} (e^{2\pi i\nu} - 1) \frac{\sin(x(p-q))}{p-q} \right)$$

- ▶ It is challenging to do microscopic

- ▶ Overlaps are too small $\sim e^{-cN}$
- ▶ Too many soft modes $\sim e^{cN}$



- ▶ Heuristic approach instead!

Dressing: inhomogeneous and complex valued!!!

$$\frac{n_F(q)}{\pi} (e^{2\pi i\nu} - 1) = \frac{e^{2\pi i\nu_T(q)} - 1}{\pi}, \quad \nu \rightarrow \nu_T(q) = \frac{1}{2\pi i} \log(1 + (e^{2\pi i\nu} - 1)n_F(q))$$




$$\tau(x) \approx \exp \left(-ix \int_{-\infty}^{\infty} \nu_T(q) dq + \int_{\mathbb{R}^2} \frac{\nu_T(q)\nu_T(q')}{(q' - q + i0)^2} dqdq' \right)$$



$$\text{Tr}(e^{-\beta H} \mathcal{O}(x, t) \dots) / \text{Tr} e^{-\beta H} = \langle \mathcal{O}(z = x + it) \dots \rangle_{\mathbb{S}^1 \times \mathbb{R}^1} \stackrel{z \rightarrow z' = e^{2\pi z / \beta}}{=} \sim \langle \mathcal{O}(z') \dots \rangle_{\mathbb{R}^2}$$

CFT prediction for correlation length:

$$\tau(x) \Big|_{T=0} = \frac{\mathcal{A}}{x^{\nu^2}} \implies \tau(x) = \frac{\mathcal{A}}{(\sinh(xT)/T)^{\nu^2}} \sim e^{-x/\xi} \implies 1/\xi \sim T^{??}$$



Generalized Sine Kernels

$$\tau(x) = \det_{[-\pi, \pi]} \left(1 + \hat{V} + \delta \hat{V} \right) - \det_{[-\pi, \pi]} \left(1 + \hat{V} \right)$$

XY model [A.G. Izergin, V.S. Kapitonov, N.A. Kitanine, solv-int/9710028]

$$V(p, q) = -\frac{\omega_F(q)}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}}, \quad \delta V(p, q) = -\frac{\omega_F(q)}{\pi} e^{-i(p+q)x/2} e^{-\frac{i(p-q)}{2}}$$

$$\omega_F(q) = \frac{1}{2} \left(1 - e^{i\theta(q)} \tanh \frac{\beta E(q)}{2} \right)$$

$$\mathbf{H}_{XY} = -\frac{1}{2} \sum_{j=1}^L \left[\frac{1+\gamma}{2} \sigma_j^x \sigma_{j+1}^x + \frac{1-\gamma}{2} \sigma_j^y \sigma_{j+1}^y + h \sigma_j^z \right], \quad \tau(x) \equiv \frac{\text{Tr} \sigma_{x+1}^x \sigma_1^x e^{-\beta \mathbf{H}_{XY}}}{\text{Tr} e^{-\beta \mathbf{H}_{XY}}}$$

Spectrum of fermionic (Majorana) excitations

$$E(q) = \sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}$$

Bogolyubov rotation angle

$$e^{i\theta(q)} = \frac{h - \cos q - i\gamma \sin q}{\sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}}$$

Form-factors

$$\tau(x) = \sum_{\mathbf{q}} |\langle \mathbf{k} | \mathbf{q} \rangle|^2 e^{-ix \left(\sum_{i=1}^{N+1} k_i - \sum_{i=1}^N q_i \right)}$$

with

$$e^{ikL} = e^{-2\pi i \nu(k)}, \quad e^{iqL} = 1.$$



$$|\langle \mathbf{k} | \mathbf{q} \rangle|^2 = A \left(\prod_{i=1}^{N+1} \frac{\sin \pi \nu(k_i)}{L} \right)^2 \frac{\prod_{i>j}^{N+1} \sin^2 \frac{k_i - k_j}{2} \prod_{i>j}^N \sin^2 \frac{q_j - q_i}{2}}{\prod_{i=1}^{N+1} \prod_{j=1}^N \sin^2 \frac{k_i - q_j}{2}}$$

$$\tau(x) = \det_{[-\pi, \pi]} \left(1 + \hat{V} + \delta \hat{V} \right) - \det_{[-\pi, \pi]} \left(1 + \hat{V} \right)$$

$$V(p, q) = -\frac{e^{2\pi i \nu(q)} - 1}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} + O(e^{-\#x}), \quad \delta V(p, q) = -\frac{e^{2\pi i \nu(q)} - 1}{\pi} e^{\frac{-i(p+q)x}{2}} e^{\frac{-i(p-q)}{2}}$$

$$e^{2\pi i\nu(k)} = 1 - 2\omega_F(k) = e^{i\theta(k)} \tanh \frac{\beta E(k)}{2}$$

$$\nu(\pi) - \nu(-\pi) = \delta \in \mathbb{Z}$$



$$\delta = 1$$

Total number of solutions

$$e^{iqL} = 1, \quad q_j = \frac{2\pi}{L} \left(-\frac{L+1}{2} + j \right), \quad j = 1, 2, \dots, L$$

$$e^{ikL} = e^{-2\pi i\nu(k)}, \quad k_j \approx \frac{2\pi}{L} \left(-\frac{L+1}{2} + j - \nu_j \right), \quad j = 1, 2, \dots, L+\delta$$

$$\nu(k) \rightarrow \nu_\delta(k) = \nu(k) - \delta \frac{k + \pi}{2\pi} \implies e^{ik(L+\delta)} = (-1)^\delta e^{-2\pi i\nu_\delta(k)}$$

For $\delta = 1$ there is only **one!** form-factor

$$\tau(x) = \sum_{q_1 < \dots < q_L} |\langle \mathbf{k} | \mathbf{q} \rangle|^2 e^{-ix \left(\sum_{i=1}^{L+1} k_i - \sum_{i=1}^L q_i \right)} = |\langle \mathbf{k} | \mathbf{q} \rangle|^2 e^{-ix \left(\sum_{i=1}^{L+1} k_i - \sum_{i=1}^L q_i \right)}$$

$$\Delta P \equiv \sum_{i=1}^{L+1} k_i - \sum_{i=1}^L q_i \approx \pi - \int_{-\pi}^{\pi} \nu(q) dq.$$

$$\tau(x) = \exp \left(-i\pi x + ix \int_{-\pi}^{\pi} \nu(q) dq - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu(q) - \nu(k) - (q-k)/2\pi}{2 \sin \frac{q-k}{2}} \right]^2 \right).$$

$$\delta = 0$$

$$\mathbf{q}^{(a)} = \{q_1, \dots, q_{a-1}, q_{a+1}, \dots, q_L\}, \quad a = 1, 2, \dots, L.$$

For $a \sim L$, $a \sim L - a$

$$|\langle \mathbf{k} | \mathbf{q}^{(a)} \rangle|^2 = \frac{e^{-2\pi i \nu(q_a)} - 1}{L} \exp \left(-\frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu(q) - \nu(k)}{2 \sin \frac{q-k}{2}} \right]^2 - \int_{-\pi}^{\pi} \nu(q) \cot \frac{q - q_a + i0}{2} dq \right)$$

$$\tau(x) = e^{-ix \sum_{j=1}^L (k_j - q_j)} \sum_{a=1}^L |\langle \mathbf{k} | \mathbf{q}^{(a)} \rangle|^2 e^{-ix q_a} = T_0(x) Y_0(x)$$

with

$$T_0(x) = \exp \left(ix \int_{-\pi}^{\pi} \nu(q) dq - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu(q) - \nu(k)}{2 \sin \frac{q-k}{2}} \right]^2 \right).$$

and

$$Y_0(x) = \int_{-\pi}^{\pi} \frac{dk}{2\pi} (e^{-2\pi i \nu(k)} - 1) e^{-ikx} \exp \left(-\int_{-\pi}^{\pi} \nu(q) \cot \frac{q - k + i0}{2} dq \right).$$

Same approach for $\delta = 1$???

$$\delta < 0$$

$$\mathbf{q}^{a_1, \dots, a_n} = \{q_1, \dots, \hat{q}_{a_1}, \dots, \hat{q}_{a_n}, \dots, q_L\} \quad \delta = 1 - n$$

$$\Delta P_{a_1, \dots, a_n} = \sum_{i=1}^{L-n+1} k_i - \sum_{i=1}^L q_i + \sum_{i=1}^n q_{a_i} \approx \delta\pi - \int_{-\pi}^{\pi} \nu(q) dq + \sum_{i=1}^n q_{a_i}.$$

$$e^{-ix\Delta P_{a_1, \dots, a_n}} |\langle \mathbf{k} | \mathbf{q}^{a_1, \dots, a_n} \rangle|^2 = \mathcal{A}_\delta[\nu] \prod_{i>j}^n \left(2 \sin \frac{q_{a_i} - q_{a_j}}{2} \right)^2 \prod_{i=1}^n \mathcal{Y}_{a_i},$$

$$\tau(x) = \det_{1 \leq j, k \leq n} [Y_\delta(x + j - k)] \exp \left(ix \int_{-\pi}^{\pi} \nu_\delta(q) dq - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu_\delta(q) - \nu_\delta(k)}{2 \sin \frac{q-k}{2}} \right]^2 \right),$$

where $\nu_\delta(q) \equiv \nu(q) - \delta(q + \pi)/(2\pi)$ has zero winding number and $Y_\delta(x)$ stands for

$$Y_\delta(x) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \left(e^{-2\pi i \nu(q)} - 1 \right) \exp \left(-i(x - \delta)q + i\delta\pi - \int_{-\pi}^{\pi} dk \nu_\delta(k) \cot \frac{q - k + i0}{2} \right).$$

$$e^{2\pi i\nu(k)} = 1 - 2\omega_F(k) = e^{i\theta(k)} \tanh \frac{\beta E(k)}{2} \quad \tau(x) = \mathcal{A}(T, h, \gamma) e^{-x/\xi(T, h, \gamma)}$$

Ferromagnetic $h \leq 1$ ($\delta = 1$)

$$\tau(x) = \exp \left(-i\pi x + ix \int_{-\pi}^{\pi} \nu(q) dq - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu(q) - \nu(k) - (q-k)/2\pi}{2 \sin \frac{q-k}{2}} \right]^2 \right)$$

$$\xi^{-1} = -i \int_{-\pi}^{\pi} \nu(q) dq = -\frac{1}{2\pi} \int_{-\pi}^{\pi} dk \log \tanh \frac{\beta E(k)}{2}$$



Paramagnetic $h > 1$ ($\delta = 0$)

$$\tau(x) = T_0(x) \int_{-\pi}^{\pi} \frac{dk}{2\pi} (e^{-2\pi i\nu(k)} - 1) e^{-ikx} \exp \left(- \int_{-\pi}^{\pi} \nu(q) \cot \frac{q-k+i0}{2} dq \right)$$



$$\xi^{-1} = -\frac{1}{2\pi} \int_{-\pi}^{\pi} dk \log \tanh \frac{\beta E(k)}{2} + \log y_{\pm}, \quad y_{\pm} = \frac{h + \sqrt{h^2 + \gamma^2 - 1}}{1 \pm \gamma}$$

$$\log \mathcal{A} = \log \frac{2}{\beta \sqrt{h^2 + \gamma^2 - 1}} - i \int_{-\pi}^{\pi} dq \nu(q) \frac{e^{iq} + y_{+}}{e^{iq} - y_{+}} - \frac{1}{4} \int_{-\pi}^{\pi} dq \int_{-\pi}^q dk \frac{(\nu(q) - \nu(k))^2}{\sin^2 \frac{q-k}{2}}$$

$$\beta = 1.1, \gamma = 0.25$$



Correlation function



$$\mathcal{A} = XY,$$

where:

$$X = \prod_{l=1}^{\infty} \frac{(1 - \lambda_1^{-1} f_{2l-1}) (1 - \lambda_1^{-1} g_{2l-1}) (1 - \lambda_2^{-1} f_{2l-1}) (1 - \lambda_2^{-1} g_{2l-1})}{(1 - \lambda_1^{-1} f_{2l}) (1 - \lambda_1^{-1} g_{2l}) (1 - \lambda_2^{-1} f_{2l}) (1 - \lambda_2^{-1} g_{2l})}$$

$$Y = \prod_{i,j=1}^{\infty} \frac{(1 - f_{2j} f_{2i-1})(1 - f_{2i} f_{2j-1})(1 - g_{2j} g_{2i-1})(1 - g_{2i} g_{2j-1})}{(1 - f_{2j} f_{2i})(1 - f_{2j-1} f_{2i-1})(1 - g_{2j} g_{2i})(1 - g_{2j-1} g_{2i-1})} \times \\ \times \frac{(1 - f_{2j} g_{2i-1})(1 - f_{2i} g_{2j-1})(1 - g_{2j} f_{2i-1})(1 - g_{2i} f_{2j-1})}{(1 - f_{2j} g_{2i})(1 - g_{2j} f_{2i})(1 - g_{2j-1} f_{2i-1})(1 - f_{2j-1} g_{2i-1})}$$

and $\lambda_1, \lambda_2, f, g$ are defined as

$$\lambda_1 = \left\{ h + [h^2 - (1 - \gamma^2)]^{1/2} \right\} / (1 - \gamma), \quad \lambda_2 = \left\{ h - [h^2 - (1 - \gamma^2)]^{1/2} \right\} / (1 - \gamma)$$

$$f_k = \frac{h + W_k}{1 - \gamma^2} - \left[\left(\frac{h + W_k}{1 - \gamma^2} \right)^2 - 1 \right]^{1/2}, \quad g_k = \frac{h - W_k}{1 - \gamma^2} - \left[\left(\frac{h - W_k}{1 - \gamma^2} \right)^2 - 1 \right]^{1/2}$$

with

$$W_k = \left\{ \gamma^2 h^2 - (1 - \gamma^2) [\gamma^2 + (k\pi)^2 \beta^{-2}] \right\}^{1/2}$$

Fredholm to Toeplitz1

$$\frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} \sim \sum_{n=0}^{x-1} e^{in(q-p)} \rightarrow \sum_{n=0}^{x-1} a_n(p) e^{in(q-p)}.$$

$$\widehat{S}_\nu(p, q) = \frac{e^{2\pi i \nu(p)} - 1}{2\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} \sim \sum_n \mathcal{A}_{qn} \mathcal{B}_{np}$$

$$\mathcal{A}_{qn} = e^{iqn}, \quad \mathcal{B}_{np} = \frac{e^{2\pi i \nu(p)} - 1}{2\pi} a_n(p) e^{-inp}.$$

$$\det(1 + \mathcal{A}\mathcal{B}) = \det(1 + \mathcal{B}\mathcal{A})$$

$$\det(1 + \widehat{S}^a) = \det_{0 \leq n, m \leq x-1} (\delta_{nm} + T_{nm}), \quad T_{nm} = \int_{-\pi}^{\pi} \frac{dq}{2\pi} a_n(q) (e^{2\pi i \nu(q)} - 1) e^{-i(n-m)q}.$$

For $a_n = 1$ the matrix T_{nm} transforms into the Toeplitz one

$$\det(1 + \widehat{S}^a) = \det_{0 \leq n, m \leq x-1} c_{n-m}, \quad c_k = \int_{-\pi}^{\pi} \frac{dq}{2\pi} e^{2\pi i \nu(q)} e^{-ikq}.$$

Fredholm to Toeplitz2

$$\det \left(1 + \hat{S}_\nu + \delta \hat{V}_\nu \right) - \det \left(1 + \hat{S}_\nu \right) = \frac{\partial}{\partial \alpha} \det \left(1 + \hat{S}_\nu + \alpha \delta \hat{V}_\nu \right) \Big|_{\alpha=0}.$$

$$\delta V_\nu(p, q) = -\frac{e^{2\pi i \nu(p)} - 1}{2\pi} e^{-i(x+1)p/2} e^{-i(x-1)q/2}$$

We choose $a_0(q) = 1 - \alpha e^{-ixq}$ and $a_n(q) = 1$ for $n \geq 1$

$$\det \left(1 + \hat{S}_\nu + \alpha \delta \hat{V}_\nu \right) = \det \left(1 + \widehat{S}^a \right) = \det \begin{pmatrix} c_0 - \alpha c_x & c_{-1} - \alpha c_{x-1} & \dots & c_{-x+1} - \alpha c_1 \\ c_1 & c_0 & \dots & c_{-x+2} \\ \cdot & \cdot & \ddots & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ c_{x-1} & c_{x-2} & \dots & c_0 \end{pmatrix}.$$

$$\frac{\partial}{\partial \alpha} \det \left(1 + \hat{S}_\nu + \alpha \delta \hat{V}_\nu \right) \Big|_{\alpha=0} = (-1)^x \det \begin{pmatrix} c_1 & c_0 & \dots & c_{-x+2} \\ c_2 & c_1 & \dots & c_{-x+3} \\ \cdot & \cdot & \ddots & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ c_x & c_{x-1} & \dots & c_1 \end{pmatrix} = \det_{0 \leq n, m \leq x-1} \tilde{c}_{n-m},$$

Fredholm to Toeplitz3

$$\det \left(1 + \hat{S}_\nu + \delta \hat{V}_\nu \right) - \det \left(1 + \hat{S}_\nu \right) = \det_{0 \leq n, m \leq x-1} \tilde{c}_{n-m}$$

$$\tilde{c}_k = - \int_{-\pi}^{\pi} \frac{dq}{2\pi} e^{2\pi i \nu(q)} e^{-i(k+1)q} = \int_{-\pi}^{\pi} \frac{dq}{2\pi} e^{2\pi i \nu_1(q)} e^{-ikq}.$$

$$\nu_1(q) = \nu(q) - \frac{q + \pi}{2\pi}$$

$$\det \left(1 + \hat{S}_\nu + \delta \hat{V}_\nu \right) - \det \left(1 + \hat{S}_\nu \right) = \det \left(1 + \hat{S}_{\nu_1} \right)$$

Szegő theorem for Toeplitz determinant [Szegő (1915), Fisher & Hartwig (1969)]

$$\log \det_{0 \leq i, j \leq x-1} c_{i-j} = x k_0 + \sum_{n=1}^{\infty} n k_n k_{-n}$$

$$\nu_\delta(q) = \frac{-1}{2\pi i} \sum_{n=-\infty}^{\infty} k_n e^{iqn}$$

$$-\frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dp \left[\frac{\nu_\delta(q) - \nu_\delta(p)}{2 \sin \frac{q-p}{2}} \right]^2 = \sum_{n=1}^{\infty} n k_n k_{-n},$$

Field theory treatment = Microscopic bosonization ($T = 0$)

- ▶ Form-Factor summation


$$\tau(x, t) = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}}} = \det(1 + \hat{V})$$

- ▶ Orthogonality Catastrophe: $|\langle \mathbf{q} | \mathcal{O} | \mathbf{k}_{\text{vac}} \rangle|^2 = \mathcal{A}/N^{2\alpha}$
- ▶ Soft-mode summation

$$\tau(x, t) \sim \sum_{\text{IR}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}}} = \langle e^{\sqrt{\alpha}\varphi(x,t)} e^{-\sqrt{\alpha}\varphi(0,0)} \rangle = \frac{\mathcal{A}}{(x - k_F t)^\alpha (x + k_F t)^\alpha}$$

- ▶ Nonlinear contributions

$$\tau(x, t) \sim \sum_{Q+\text{IR}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}} + ix(Q - k_F) + it(Q^2 - k_F^2)/2} = \frac{\mathcal{B}}{\sqrt{t}(x - k_F t)^{\tilde{\alpha}}(x + k_F t)^\alpha}$$






Slavnov (1989); Slavnov and Korepin (1991); A. Shashi, L. I. Glazman, J.-S. Caux, and A. Imambekov (2011); N. Kitanine, K.K. Kozłowski, J.-M. Maillet, N.A. Slavnov, and V. Terras (2009-2012); K.K. Kozłowski, J.-M. Maillet (2015);

Dynamics (Preliminary) ($T = 0$)

$$\tau(x, t) = \frac{G^2(1-\nu)G^2(1+\nu)}{(2i(t-x))\nu^2(2i(x+t))\nu^2} e^{-2i\nu x} +$$

$$\frac{G^2(1-\nu)G^2(\nu)}{\nu^2(2i(t-x))(1+\nu)^2(2i(x+t))\nu^2} \left(\frac{x-t}{x+t}\right)^{2\nu} \frac{e^{-i(t-x)^2/(2t)-2i\nu x}}{(x/t-1)^2} \sqrt{\frac{2\pi}{-it}} \theta(x^2 > t^2) +$$

$$+ \frac{G^2(-\nu)G^2(1+\nu)}{\nu^2(2i(t-x))(1-\nu)^2(2i(x+t))\nu^2} \left(\frac{x+t}{x-t}\right)^{2\nu} \frac{e^{i(t-x)^2/(2t)-2i\nu x}}{(x/t-1)^2} \sqrt{\frac{2\pi}{it}} \theta(x^2 < t^2) + (\nu \rightarrow \nu + \mathbb{Z})$$



Dynamics (Preliminary)

$$\nu_T(q) = \frac{1}{2\pi i} \log \left[1 + (e^{2\pi i \nu} - 1) n_F(q) \right] \theta(x - qt) - \frac{1}{2\pi i} \log \left[1 + (e^{-2\pi i \nu} - 1) n_F(q) \right] \theta(qt - x).$$

$$\text{Erf} \left(\frac{(x - qt)(1 + i)}{2\sqrt{t}} \right) \rightarrow \text{Sign}(x - qt)$$




Figure: The exact expressions (solid lines) vs. the asymptotic (dotted lines) for $\nu = 0.4$. (Real part is black and absolute value is red).

Summary and outlook

- ▶ Riemann Hilbert Problem for Fredholm determinant
- ▶ Phase shift dressing
- ▶ Different types of soft mode contributions
- ▶ Universality
- ▶ Asymptotic for classical integrable models?
- ▶ Relation with QTM? Thermal form-factors?

Extra slides

Soft mode summation



$$\frac{|\langle P_k, H_k | c_p | \mathbf{k} \rangle_Q|^2}{|\langle \text{vac} | c_p | \mathbf{k} \rangle_Q|^2} = \left(\det_{1 \leq i, j \leq k} \frac{1}{p_i + h_j - 1} \right)^2 \left(\frac{\sin \pi F_+}{\pi} \right)^{2k} \prod_{j=1}^k \frac{\Gamma(p_j - F_+)^2}{\Gamma(p_j)^2} \prod_{j=1}^k \frac{\Gamma(h_j + F_+)^2}{\Gamma(h_j)^2}$$

$$P_q - Q = \int_{-1}^1 dq F(q) + \frac{2\pi}{L} \sum_{j=1}^k (p_j + h_j - 1)$$

Auxiliary Fermions

$$\{\psi_n, \psi_m^+\} = \delta_{nm}, \quad \{\psi_n, \psi_m\} = \{\psi_n^+, \psi_m^+\} = 0.$$

$$\psi_n |0\rangle = 0, \quad \text{if } n > 0, \quad \psi_n^+ |0\rangle = 0, \quad \text{if } n \leq 0$$

$$|P_k, H_k\rangle = \psi_{p_1}^+ \dots \psi_{p_k}^+ \psi_{1-q_1} \dots \psi_{1-q_k} |0\rangle$$

Bosonization

$$\psi^+(z) = \sum_{n \in \mathbb{Z}} z^n \psi_n^+, \quad \psi(z) = \sum_{n \in \mathbb{Z}} z^{-n} \psi_n.$$

$$J(z) =: \psi^+(z)\psi(z) := \sum_{k \in \mathbb{Z}} \frac{\sum_{j \in \mathbb{Z}} \psi_j^+ \psi_{j+k}}{z^k} \equiv \sum_{k \in \mathbb{Z}} \frac{J_k}{z^k},$$

$$z \frac{\partial \varphi(z)}{\partial z} = J(z), \quad \varphi(z) = \varphi_-(z) - \varphi_+(z) + J_0 \log z + N_0$$

$$\varphi_+(z) = \sum_{k > 0} \frac{J_k}{k z^k}, \quad \varphi_-(z) = \sum_{k > 0} z^k \frac{J_{-k}}{k}$$

$$e^{iy(P_q - Q)} \frac{|\langle P_k, H_k | c_p | \mathbf{k} \rangle_Q|^2}{|\langle \text{vac} | c_p | \mathbf{k} \rangle_Q|^2} = e^{iy \Delta P} \langle 0 | e^{F_+ \varphi_+(1)} | P_k, H_k \rangle \langle P_k, H_k | e^{F_+ \varphi_-(z)} | 0 \rangle$$

$$\Delta P = - \int_{-1}^1 dq F(q), \quad z = e^{2\pi i y / L}$$

Bosonization

$$\begin{aligned} \sum_{P_k, H_k} e^{iy(P_q - Q)} \frac{|\langle P_k, H_k | c_p | \mathbf{k} \rangle_Q|^2}{|\langle \text{vac} | c_p | \mathbf{k} \rangle_Q|^2} &= e^{iy\Delta P} \sum_{P_k, H_k} \langle 0 | e^{F_+ \varphi_+(1)} | P_k, H_k \rangle \langle P_k, H_k | e^{F_+ \varphi_-(z)} | 0 \rangle = \\ &= e^{iy\Delta P} \langle 0 | e^{F_+ \varphi_+(1)} e^{F_+ \varphi_-(z)} | 0 \rangle = e^{iy\Delta P} \langle 0 | e^{F_+^2 [\varphi_+(1), \varphi_-(z)]} | 0 \rangle = \frac{e^{iy\Delta P}}{(1-z)^{F_+^2}} \end{aligned}$$

$$\sum_{P_k, H_k} e^{iy(P_q - Q)} |\langle P_k, H_k | c_p | \mathbf{k} \rangle_Q|^2 = |\langle \text{vac} | c_p | \mathbf{k} \rangle_Q|^2 \frac{e^{iy\Delta P}}{(1-z)^{F_+^2}} = \frac{\mathcal{A}}{N^{F_+^2}} \frac{e^{iy\Delta P}}{(1 - e^{2\pi i/L})^{F_+^2}}$$

$$\sum_{P_k, H_k} e^{iy(P_q - Q)} |\langle P_k, H_k | c_p | \mathbf{k} \rangle_Q|^2 = \frac{\mathcal{A} e^{iy\Delta P}}{(-2k_F i y)^{F_+^2}}$$

- ▶ Any soft modes excitations
- ▶ Non-linear bosonization